

Contents lists available at [SciVerse ScienceDirect](http://www.sciencedirect.com)

Tzu Chi Medical Journal

journal homepage: www.tzuchimedjnl.com

Clinical Practice and Therapeutics

Genetic counseling on amniocyte level II mosaicism

Chun-Ying Weng^a, Shao-Yin Chu^{a,b,d,*}, Tzu-Ying Li^a, Jye-Siung Fang^c, Ming-Liang Lee^{a,d}^a Department of Pediatrics, Buddhist Tzu Chi General Hospital, Hualien, Taiwan^b Department of Medical Education, Buddhist Tzu Chi General Hospital, Hualien, Taiwan^c Department of Molecular Biology and Human Genetics, Tzu Chi University, Hualien, Taiwan^d Department of Medicine, Tzu Chi University, Hualien, Taiwan

ARTICLE INFO

Article history:

Received 25 May 2011

Received in revised form

24 June 2011

Accepted 30 June 2011

A 36-year-old woman who was a junior college graduate and a housewife received amniocentesis because of her advanced maternal age. Giemsa banding revealed the karyotype of the amniocytes was 46, XY. More than two cells with 46, XY, t (1; 2) (p21; q22) in one colony, however, were found among the 15 colonies analyzed (Fig. 1). Mosaicism level II was diagnosed.

Genetic counseling was arranged because of the parents' concerns. The counseling process included information on chromosomes and karyotypes. Mosaicism is the coexistence of a normal and abnormal karyotype. Mosaicism detected prenatally is categorized into three levels. Level I is when a single abnormal cell is found in one colony, and is considered an artifact of the culture. If more than two cells with the same chromosome abnormality are found in one colony, or the same abnormality is found in more than two colonies from one culture, and the abnormality is not seen in colonies from other independent cultures, it is considered level II mosaicism (Fig. 1). These cases are usually found to be pseudo-mosaicism, and additional studies may be performed. Level III mosaicism is defined as the detection of two or more cells with the same chromosome abnormality distributed over two or more independent cultures. Those cases are likely to be true mosaicism.

According to the standard procedures in the cytogenetic laboratory, additional studies were not needed to eliminate the possibility of chromosomal abnormality in our case. The parents were counseled that major malformations occur in 3–4% of live births. Prenatal genetic counseling was also recommended for the next pregnancy.

The follow-up of this pregnancy was uneventful, and the baby was born at full term with a birth body weight of 2870 gm. The perinatal course was smooth and repeat blood chromosome testing revealed a karyotype of 46, XY (20 cells).

Further reading

- [1] Shaffer LG, Slovak ML, Campbell LC. International Standing Committee on Human Cytogenetic Nomenclature (ISCN): An International System for human cytogenetic nomenclature. KARGER; 2009. p. 77.
- [2] Canadian College of Medical Geneticists (CCMG) Cytogenetic Guidelines. Canada: CCMG; 2003. pp. 2–6.
- [3] Dimaio MS, Fox JE, Mahoney MJ. Cytogenetic Abnormalities, chapter 1, Chromosomal mosaicism -prenatal diagnosis, Prenatal diagnosis: cases and clinical challenges. 1st ed. UK: Blackwell Publishing Ltd; 2010. pp. 11–13.

* Corresponding author. Department of Pediatrics, Buddhist Tzu Chi General Hospital, 707, Section 3, Chung-Yang Road, Hualien, Taiwan. Tel.: +886 38561825x3649; fax: +886 38563532.

E-mail address: chu_chu@tzuchi.com.tw (S.-Y. Chu).




Fig. 1. (A) There are more than two cells with the same abnormality in one colony among the 15 colonies analyzed on culture slide number 2. (B) The karyotype of the green cells is 46, XY. (C) The karyotype of the lined cells is 46, XY/46, XY, t (1; 2) (p21; q22). No other abnormal cells can be seen in colonies from slide 1 and slide 3.